

ASSURING THE HEALTH AND WELL-BEING OF THE PREGNANT MARE.

We often think of pregnancy as a delicate and fragile condition. When it comes to horses, the perception is perhaps due to the mare's relatively poor reproductive performance in comparison to other domestic animals. However, in a natural setting, the mare does comparatively well reproductively. Therefore, this seemingly poor performance is due as much to improper management as to any reproductive deficiency. Fortunately, management is something we can control.

As a conscientious owner, you probably have many questions about caring for your expectant mare. In truth, you may be a little worried. Relax. With a little TLC, your mare should progress through her pregnancy without mishap. Proper nutrition, deworming, exercising, and vaccinations will help ensure a healthy pregnancy, and you can look forward to the birth of your foal with greater confidence.

PRECARIOUS BEGINNINGS

The earliest days of an embryo's existence are perhaps the most precarious. During the first 30 days, there is a 10-15% chance that the embryo will be resorbed. Stress, illness, uterine infection, hormonal abnormalities, the presence of twins, and other factors have been implicated in early embryonic loss. Often, the cause remains undetermined.

When the mare conceives, the fertilized egg (zygote) travels down the fallopian tubes and enters the uterus around day 6-7. It migrates throughout the uterus until about day 16 and typically implants into the uterine wall at 6-8 weeks. By day 12-13, the embryonic vesicle is usually large enough to be detected by ultrasonic examinations, during which an image is made by bouncing sound waves off tissues. For practical reasons, some breeding farms simply tease the mare 14-20 days after her last breeding date to see if she comes back into estrus (heat). If she does not, the pregnancy may then be confirmed by ultrasound or trans-rectal palpation at approximately 30-35 days post-breeding.

Neither teasing, palpation, nor ultrasound has been shown to harm the developing embryo or endanger the pregnancy. However, because of the embryo's uncertain beginning, it may be wise to have the pregnancy reconfirmed at 45, 60, or 90 days.

TROUBLESOME TWINS

Some reproductive specialists recommend an ultrasound exam at 14-16 days post-ovulation to detect twins. Early detection of twins provides an opportunity to eliminate one embryo, thus allowing the other to develop normally. This is commonly done because twins pose a number of risks:

- In 95% of mares with twin embryos, one or both embryos are resorbed during the first 60 days. However, waiting to see if this occurs naturally could delay or interfere with a subsequent successful pregnancy.
- Of the small percentage of twins that survive in utero past 50 days, it is highly unlikely that two healthy will be born. If either survives, it may be small and weak.
- Most twins surviving past 50 days will spontaneously abort 6-8 months.
- Mares carrying twins are more likely to give birth prematurely (before 300-320 days). Premature foals may have serious medical problems and are less likely to survive.

HELPING NATURE TAKE ITS COURSE

Good broodmare management is the best aid for helping the mare make it through the critical first 30-60 days of pregnancy.

The mare should go into the breeding season fit and perhaps gaining weight. Severely underweight mares will have more trouble conceiving than will mares of appropriate weight.

Avoid stressing the mare as much as possible. Stress can cause a drop in progesterone, a hormone which helps maintain pregnancy. Illness and/or fever can cause the mare's system to secrete prostaglandins, which may cause abortion.

USE GOOD JUDGMENT

- Transport your mare only if necessary.
- Use caution when exposing your mare to other horses. You should avoid any undue risk of injury or disease transmission.
- Provide nutritious forage, but don't overfeed. Supplementing with vitamins and minerals is unnecessary in mares being fed a balanced diet.
- Make sure the mare is current on vaccines and deworming. Consult your veterinarian for recommendations regarding specific vaccinations and deworming interval during pregnancy.
- Do not administer hormones or other drugs unless specifically prescribed by your equine practitioner.
- Carefully evaluate the mare before deciding whether to breed on foal heat. Consult your veterinarian.

COASTING THROUGH MIDDLE PREGNANCY

Unless there are special circumstances, during the first 7 months of pregnancy, treat your mare as you would a non-pregnant one. She will benefit from moderate riding or exercise. The ration should be composed primarily of high-quality forage in approximately the same as pre-pregnancy amounts. Extremes in weather can alter her nutritional requirements and should be taken into account when formulating the ration. She should always have plenty of clean, fresh water. The mare will also benefit from routine hoof and dental care, standard vaccinations, and regular deworming.

VACCINES

Vaccinations should be current, since infectious diseases can trigger abortions. A four-way inoculation for Eastern and Western encephalomyelitis, influenza, and tetanus is recommended at the beginning of pregnancy. A booster should be given one month prior to foaling to increase the antibody level in the mare's colostrums (first milk) and help protect the newborn foal from disease. Also, the mare should be vaccinated for equine rhinopneumonitis (commonly called virus abortion or rhino) at five, seven, and nine month's gestation. Consult with your local veterinarian regarding other vaccines that may be advisable in your area, such as rabies and botulism.

DEWORMING

Most deworming agents available today are relatively safe for pregnant mares. Consult your veterinarian to establish an effective and safe deworming schedule for your mare.

It is especially important to deworm the mare within several weeks of foaling, because the mare will be the primary source for infecting her foal with parasites. Of course, manure should always be properly disposed of.

CHANGING NEEDS

During the last four months of pregnancy, the foal will grow rapidly. To accommodate this growth, the mare's energy needs will increase. Even so, special nutritional supplements are probably unnecessary. Good-quality hay and forage should remain the bulk of the expectant mare's diet. Concentrated feeds, such as grains, may be added to the ration to bolster energy intake without adding excess bulk.

Use the mare's body conditions as your guide to how she's faring. Adjust the ration accordingly. The mare should not become obese. Your veterinarian can advise you regarding a proper nutritional program for your mare.

Exercise during the last four months of the mare's pregnancy should be light to moderate. In fact, a pastured mare will get as much exercise as she needs just grazing. Vigorous exercise is not recommended.

HOME STRETCH

The average length pregnancy in the mare is 338-343 days. However, normal gestation can range from 320-380 days. You needn't become overly concerned if your mare is past due. Prolonged gestation is not generally associated with problems or extra large foals. If your mare's pregnancy extends much past 340 days or you're concerned. Ask your veterinarian to examine her to determine if the mare is still pregnant and confirm that all is well.

SUSPECTED ABORTION

Mares do occasionally abort. If you notice a vaginal discharge or dripping milk during pregnancy, contact your veterinarian. If you find the remains of a placenta or fetus, save it for your veterinarian to examine. It may be possible to ascertain the cause of abortion and treat the mare accordingly. Mares can and do abort without ill effects. However, it's always a good idea to have her checked by your veterinarian, because some complications of abortions, such as a retained placenta, can be life-threatening to your horse.

IMPENDING BIRTH

There are obvious as well as subtle signs of impending birth. The time frame during which they occur varies from mare to mare. The most obvious and reliable are:

- Filling of the udder (two to four weeks pre-foaling)
- Distension of the teats (four to six days pre-foaling)
- Waxing of the teats (one to four days pre-foaling)
- Obvious dripping of milk

More subtle signs include:

- Softening and flattening of the muscles in the croup
- Relaxation of the vulva
- Visible changes in the position of the foal

PREPARING FOR BIRTH

Your eleven-month waiting game will be over before you know it. Ask your veterinarian any questions or concerns you have may have about caring for your expectant mare.